



# Plant Tissue Plus Soil Report

Lab No: 60001      Field: 001  
 Date Sampled: 07/11/2016      Sample: Bad (Alfalfa)

## Sufficiency Ranges


### Plant Results

	Low	Sufficient	High
N %	1.48		
P %	0.10		
K %		2.92	
Ca %	0.24		
Mg %	0.18		
S %	0.08		
Zn ppm	19		
B ppm	5.65		
Mn ppm	14		
Fe ppm		105	
Cu ppm		12	
Mo ppm			

Al ppm 85      Al and Na are  
 Na ppm 22      nonessential elements.

### Soil Results

Lab Number	178593	
pH	7.2, target = 6.8	
OM	4.5	
	Optimum	
P ppm	24	
K ppm	80	
Ca ppm		2526
Mg ppm	251	



### Key

●	Plant Result
x	Soil Result

Low (Red)  
 Sufficient (Green)  
 High (Yellow)

Lower - Upper	
N%	2.50 - 4.00
p%	0.25 - 0.45
K%	2.25 - 3.50
Ca%	0.70 - 2.50
Mg%	0.25 - 0.70
S%	0.25 - 0.50
Zn ppm	20.0 - 60.0
B ppm	25.0 - 60.0
Mn ppm	20.0 - 100.0
Fe ppm	30.0 - 250.0
Cu ppm	3.00 - 30.00

### Photos



## Yield Response Interpretation Indexes

	Almost Certain	Possible	Remote	Unlikely
DRIS	S:-21 B:-73	P:-17	N:20	K:91
PASS INI	P:-30 N:-25 B:-31 S:-38	Ca:-18 Mn:-13 Zn:-12 Mg:-28	K:-2	


**PASS DNI** K:78 P:20 N:73 B:-118 S:-53

**DRIS** Diagnosis and Recommendations Integrated System. DRIS yield response categories: *Almost Certain* index < -20, *Possible* -20 < index < -15, *Unlikely* index > 25, *Remote* any index not in any other category.

**PASS** Plant Analysis with Standardized Scores. INI = Independent Nutrient Index, DNI = Dependent Nutrient Index. PASS yield response categories: *Almost Certain* common response elements with INI < -10, *Possible* common response elements with INI+DNI < -10 and rare response elements with INI < -10, *Unlikely* any element with INI > 10, *Remote* any element not in any other category.

## Comments

- This plant sample is low or deficient in nitrogen, possibly as a result of inadequate nitrogen fertilization, excessively wet soil conditions, excessive rainfall and leaching on sandy soils, inadequate phosphorus fertilization or excessive potassium fertilization.
- Symptoms of nitrogen deficiency appear first as a light green coloring of the plant. As the deficiency becomes more severe, lower leaves turn yellow and may 'fire'.
- This plant sample is low in sulfur. This could be a result of low soil sulfur content or inadequate sulfur fertilization. Alfalfa should be topdressed with the sulfate form of sulfur. Other crops should be sidedressed with a sulfate form. All crops could be foliar treated with a liquid form of sulfur. Follow the manufacturers recommendations.
- This plant sample is low in boron. This may be a result of low soil boron availability, inadequate boron fertilization or excessive potassium fertilization. This problem is most likely to occur on sandy, alkaline or low organic matter soils especially during droughty conditions.
- Apply 2 or 3 lbs B/a once in the rotation or 0.5 to 1.0 lb B/a each year. Borax and sodium tetraborate are the most commonly used boron fertilizers. Solubor, sodium pentaborate and boric acid are used occasionally for soil application or in foliar sprays.



— 7.2 pH  
 - - - 6.8 Target pH

Soil pH is high.  
 There is no lime recommendation.

## Interpretations

Plant analysis results are interpreted by one or more of three methods: Sufficiency Range (SR); Diagnosis and Recommendation Integrated System (DRIS); or Plant Analysis with Standardized Scores system (PASS). By comparing the three methods of interpretation, it should be possible to arrive at a clearer picture of the plant's nutrient status than by using only one method.

The SR system is based on the relationship between nutrient concentration and yield. If an associated soil test shows conditions that can cause nutrient deficiency (low concentration, pH problem, etc...), then an increase in supply through corrections or additions will increase concentrations in the plant. The concentration range identified as sufficient is defined to result in 95 to 100% of maximum yield. The system is sensitive to plant maturity and plant part sampled. Interpretations are reliable only when used for the specific plant part sampled at the specific growth stages where interpretations have been developed. The diagram integrates plant analysis and the soil test results where plant analysis sufficiency range and optimum soil test levels are indicated by the middle circle. The inner circle indicates deficiency or below-optimum levels, and the outer circle marks above-optimum levels. Plant nutrient concentration survey data are substituted when sufficiency range information is not available. Survey data have not been evaluated by yield response calibrations and are to be used only for general comparison.

The DRIS is a method to evaluate various combinations of ratios of nutrient concentrations rather than the actual concentrations. These ratios are combined mathematically to give an index. An index of zero is ideal. The more negative the index, the greater the potential of nutrient deficiency; the greater the index, the greater the potential for excessive concentrations. The sum of these indices for a given analysis must be zero. To avoid errors in diagnosing deficiencies when deviations from zero are really random error, an in-balance range is defined as -15 to +25. DRIS norms are available for alfalfa, apple, corn, celery, lettuce, millet, oat, potato, grain sorghum, tomato, and wheat.

The PASS system is a hybrid that combines an Independent Index (INI) as in SR, and a Dependent Index (DNI) as in DRIS. The PASS INI section is similar to SR, but instead of a nutrient category, a continuous index based on the statistical standardized score is determined and expressed on the DRIS scale. INI values between -10 and +10 are considered sufficient. An INI value of less than -10 equals the critical level and is considered deficient. The further below -10 the INI values are, the more likely the crop will respond to increased availability of that nutrient. To avoid predicting yield responses when they are unlikely, as sometimes occurs in both SR and DRIS, the nutrients in the PASS INI system are divided into two groups: those for which yield response is common, and those for which yield response is rare. The PASS DNI is similar to DRIS where paired nutrient ratios are calculated and compared to optimum values with the standardized scores approach and expressed on the DRIS scale. The DNI is best used to confirm a deficiency indicated by the INI. Only the nutrients in the common response group of the INI are included in the DNI. PASS norms are available for alfalfa, corn, and soybean.

Source: University of Wisconsin Cooperative Extension