



## Corn silage uNDF: Potential impact factor relative to feed conversion efficiency

John Goeser for *Progressive Dairy*

*The lignified fiber measure has changed, though, thanks to a new forage measure – undigestible neutral detergent fiber (NDF) at 240 hours (uNDF240). This advance has been equivalent to adding a 3-9x40 optic on top of your favorite rifle previously outfitted with open sights.*

### AT A GLANCE

Greater uNDF240 in corn silage could decrease the milk and economic performance potential for your herd.

Lignin in corn silage and forage has been compared to the rebar in concrete, providing the structural integrity and strength to hold the road or driveway together. This added strength is great for your driveway, but added strength in forage fiber corresponds to limited digestion and energy potential per pound. Less lignin is desirable in the diet, but measuring lignin hasn't been overly useful for nutritionists. Lignin measures at the forage analysis laboratory are typically fairly small values, with limited range to the data,

thus limiting how your nutritionist can use the data on your feeds. This would be like shooting a rifle with iron sights at a target 250 yards away. The lignified fiber measure has changed, though, thanks to a new forage measure – undigestible neutral detergent fiber (NDF) at 240 hours (uNDF240). This advance has been equivalent to adding a 3-9x40 optic on top of your favorite rifle previously outfitted with open sights.

The uNDF240 measure has caught on with dairy nutritionists following Kurt Cotanch and his

colleagues' observations at the Miner Institute several years ago. More recently, Rick Grant has summarized several studies from the Miner Institute and shown that total mixed ration (TMR) uNDF240 relates to dairy cow intake and milk production potential.

Building on their findings, through unintended observations, results from a 2018 Wisconsin commercial dairy survey suggested there may be a tie between corn silage uNDF240 (percent of dry matter) and dairy performance. I presented these observations, as a case study, during my fiber-focused talk at the 2019 Mid-South Ruminant Nutrition Conference. In the survey and case study, Jared Geiser (interning with CP Feeds, Valders, Wisconsin) visited and sampled roughly 60

dairies as part of an internship project. Our aim was to investigate corn silage kernel processing score (KPS) relative to starch digestibility and dairy herd performance. Beyond KPS and starch digestibility measures, we also analyzed the corn silage samples via routine forage analysis means (NIR) for nutritive and fiber digestion parameters. The silage nutrition analyses uncovered a striking potential relationship between corn silage uNDF240 (percent of DM) and energy-corrected milk production per cow.

Before getting into uNDF240, recognize that several routine fiber digestibility parameters appeared correlated to dry matter intake (DMI), energy-corrected milk per cow (ECM) and feed conversion efficiency (pounds ECM per



### Hit it with the Haymaker!

The all-new MacDon R2 Rotary Disc Header and M1240 Windrower deliver a knock-out performance. Featuring MacDon EvenFlow™ Technology, the R2 delivers the best possible cut, conditioning, and windrow formation, allowing for quick dry down and increased productivity.

See it in action and find your local dealer at [MacDon.com](http://MacDon.com)

**MacDon**  
The Harvesting Specialists.

pound DMI). Both in vitro NDF digestibility at 30 hours (NDFD30) and the University of Wisconsin total tract NDF digestibility (TTNDFD) metrics were correlated with ECM across the roughly 60 dairies represented in the survey data. The trendlines suggested the ECM gains per unit NDFD improvement were roughly 1 and 0.8 pound, for corn silage NDFD30 and TTNDFD, respectively. These observations fall in line with Masahito Oba and Mike Allen's observations roughly 20 years ago, where the authors detailed in a well-done study that a single unit change in forage NDFD corresponds to roughly 0.4- and 0.6-pound gains in DMI and 4% fat-corrected milk, respectively. Perhaps today's cows respond to an even greater extent to improved fiber digestibility.

Getting back to corn silage uNDF240, the relationship with both DMI and ECM caught our attention as much as the NDFD relationships. Keep in mind that uNDF240 measures lignified fiber, and the rumen can't digest this fiber at all. This measure in itself may be meaningful as corn silage uNDF240 was negatively correlated (P less than 0.05; independent pairwise correlations) with DMI and ECM production. Based on the case study and survey results, a single unit of corn silage uNDF appeared to correlate with roughly 0.6 and 1.3 pounds less DMI and ECM per cow per day, respectively.

Bear in mind: These are survey data and not a controlled research study. The observations simply suggest a relationship between uNDF240 and performance may exist. However, if these observations are real, then the implications are important for your herd to grasp. Greater uNDF240 in corn silage could decrease the milk and economic performance potential for your herd.

Corn silage uNDF240 is

influenced by many factors, including environment, seed genetics, cut height, grain-to-stover ratio and agronomic management. There are numerous ways to decrease the negative potential impact. Work with your nutritionist, seed consultant and/or agronomic adviser to benchmark your current silage, and then use uNDF240 as a future silage quality indicator. Last, recognize

that uNDF240 measures lignin, and remember that lignin is important for plant health. If selecting reduced-lignin hybrids, monitor plant health closely with your agronomist during the growing season. Corn silage hygienic quality is an increasingly important area to understand. We shouldn't look to gain in fiber potential only to give up plant health in the field. ➔

*John Goeser earned a Ph.D. in animal nutrition from the University of Wisconsin – Madison, where he currently serves as an adjunct professor in the dairy science department. He also directs animal nutrition, research and innovation efforts at Rock River Lab Inc. based in Watertown, Wisconsin.*

*References omitted but are available upon request.*

## For Healthy Cows®

Time to Think Differently  
Microbiology Nutrition


# the P-One Program™

the 1st Ingredient™

The microbiome in every cow is different, yet all are fed the same TMR and expected to perform the same.

The branded strains of **Smartbacteria**, found only in the P-One Program, optimize energy transport through efficient carbohydrate metabolism.

This stabilizes and maintains rumen pH, making the P-One Program - the 1st Ingredient.



The only company bringing the fields of  
**MICROBIOLOGY & NUTRITION TOGETHER**

**Rubes®** by **Laigh Rubin**



"People, I feel we may be sending a mixed message."

 **PRIORITY IAC™**

the **Smartbacteria & Nutrition Company™**

888-444-2030 | [forhealthycows.com](http://forhealthycows.com)  
or an Authorized Reseller